

Wetland Classification as per Cowardin et al. 1979

PSS01-e0tg

Ω

PEM01-f0tg

PAB03-h0tg




Matthew J. Gray
University of Tennessee

Why Classify Wetlands?




- 1) Delineate their edges
- 2) Estimate their area
 - Boundary of development
- 3) To create maps
 - Management, Excavation, Mitigation



Caribbean

Classification of Wetland and Deepwater Habitats of the United States

<http://www.npwr.usgs.gov/resource/1998/classwet/classwet.htm>



FWS/OBS-79/31
December 1979

Lewis Cowardin (USFWS)
Virginia Carter (USGS)
Francis Golet (URI)
Edward LaRoe (NOAA)


Biological Classification System

- Wetlands
- Deepwater Habitats



Jurisdictional → USACE 1987 Manual

Boundary Between Wetland and Deepwater Systems

Non-tidal:
 >2 m (6.6 ft) in Depth
 (low water level—fall)
 Permanently flooded rivers and lakes



Tidal: Extreme low water level (spring tides)


Permanently flooded brackish marshes or marine areas

The Classification System


Hierarchical Structure

Systems (5), Subsystems (8), Classes (11),
 Subclasses (28), Dominance Type, Modifiers (3)


Marine




Estuarine




Riverine



Lacustrine




Palustrine



- Hydrologic
- Geomorphic
- Chemical
- Biological

Hierarchical Structure




SYSTEM	SUBSYSTEM	CLASS
ESTUARINE	SUBTIDAL	Brackish Bay
		Brackish Estuary
	INTERTIDAL	Brackish Bay
		Brackish Estuary
RIVERINE	TIDAL	Brackish Bay
		Brackish Estuary
	LOWER PERENNIAL	Brackish Bay
		Brackish Estuary
UPPER PERENNIAL	Brackish Bay	
	Brackish Estuary	
INTERMITTENT	Brackish Bay	
	Brackish Estuary	
LACUSTRINE	LIMNETIC	Brackish Bay
	LITTORAL	Brackish Bay
WETLANDS/DEEPWATER SYSTEMS		Brackish Bay
PALUSTRINE		Brackish Bay
		Brackish Estuary

Marine System

Open ocean overlying the continental shelf and its coastline, where salinities are >30 ppt, except at the mouths of estuaries.

1) Extreme high water
OR, limit of spring tides

2) Wetland emergent vegetation



OR,

3) Estuarine system
If #2 not present

4) Continental shelf
(ocean extent)

“High-energy Systems”

Subsystems:

Subtidal: Substrate is continuously submerged (deepwater)

Intertidal: Substrate is exposed and flooded by tides (wetland)

Marine Subsystems



Subtidal




Intertidal










Splash Zone

Estuarine System

Tidal deepwater systems and wetlands that are usually semi-enclosed by land but have open, partly-obstructed, or sporadic access to the open ocean.

1) Salinity >0.5 ppt

2) Imaginary line closes bay or river mouth



OR,

3) Wetland Vegetation
If, goes beyond #2

4) Offshore Areas
Continuously diluted w/ freshwater

“Low-energy Systems”

Subsystems:

Subtidal: Substrate is continuously submerged (deepwater)

Intertidal: Substrate is exposed and flooded by tides (wetland)

Estuarine System

Riverine System

All deepwater systems and wetlands contained within a channel **EXCEPT:**

(1) Dominated by **PERSISTENT** emergent plants, trees, shrubs, mosses, or lichens
OR,
 (2) Salinity >0.5 ppt.

Ends at Upland Boundary, Spring, or another System

Subsystems:

Lower Perennial: Low Gradient/Flow, Substrate is mud or sand.

Upper Perennial: High Gradient/Flow, Substrate is rock, gravel.



Intermittent: Water flows only during **part of the year**.

Tidal: Flow fluctuates with **tide** but freshwater.



Riverine Subsystems

Lower Perennial

Upper Perennial

Intermittent

(if non-persistent)

Lacustrine System

Freshwater deepwater habitats and wetlands with the following characteristics:

- >8 ha in Surface Area:**
- 1) Positioned in topographic depression or dammed river channel
- and,
- OR,**
- 2) There is <30% coverage of persistent emergent vegetation (non-persistent OK)
- <8 ha in Surface Area:**
- 1) #1 and 2 above (and one below)
 - 2) Wave-formed shoreline
 - 3) Deepest part of basin > 2 m depth

Subsystems:

- Limnetic:** All deepwater zones **"Ponds, Lakes & Reservoirs"**
- Littoral:** All wetlands zones (extends from the upland or Palustrine System to 2 m depth)

Lacustrine Subsystems

Size, Depth, Non-persistent

>2 m

See Handout

Palustrine System

All freshwater wetlands dominated (>30% coverage) by trees, shrubs, persistent emergents, or emergent mosses and lichens

•Non-tidal or tidal

Also, all wetlands lacking above vegetation (or dominated by non-persistent emergents) having all these 4 characteristics:

- 1) <8 ha in size
- 2) No active wave formed shoreline
- 3) Depth <2 m
- 4) Salinity <0.5ppt

No subsystem!!

Palustrine System No subsystems!!

Palustrine Illustration

Fig. 6. Ecotypifying features and examples of habitats in the Palustrine System.

Classes For Palustrine System

Describe the *general appearance* of the wetland in terms of either the dominant **vegetation** or **substrate** composition

If Horizontal Vegetative Cover (HVC) is **>30%**:

Class is distinguished based on the uppermost layer of vegetation (i.e., tallest) with HVC >30%
(e.g., 50% HVC of Trees over 60% HVC of Shrubs would be a Forested not Scrub-shrub Wetland)

If Horizontal Vegetative Cover is **<30%**:

Class is distinguished based on the texture and composition of the substrate

Classes: How many classes?
 HVC >30% (tallest strata)



Subclasses and Dominance Type

Subclass:
 Describe *finer* differences in **vegetative** life forms (often related to life history) or **substrate** characteristics (i.e., size [gravel vs. sand] or composition [sand vs. organic])


Dominance Type:
 The taxonomic category subordinate to subclass.

Use 50/20 Rule

- Dominant Plant Species (*if class is vegetation*)
- Dominant Animal Species (*if class is substrate*)

Subclasses and Dominance Type

Sampling Methods FWF 410



Types of Palustrine Classes, Subclasses, and Dominance Types


1) Rock Bottom:

- >75% HC of stones, bolders, or bedrock
- <30% HVC

Usually high-energy wetlands with well-aerated water.

- A) Bedrock: >75% bedrock substrate
- B) Rubble: <75% bedrock; >75% bedrock+bolders+stones

Dominance Types: *Ephemeralia*, *Procambarus*, *Spongilla*, and *Lymnaea* (pond snail)



Types of Palustrine Classes, Subclasses, and Dominance Types

2) Unconsolidated Bottom:

- >25% HC of soil particles smaller than stones
- <30% HVC


Usually low-energy wetlands that are flooded more permanently.


- A) Cobble-gravel: >50% c/g
- B) Sand: >50% sand
- *C) Mud: >50% silt & clay
- *D) Organic: >50% dead or live organic matter

Dominance Types: *Gammarus* (scuds), *Physa* (snail), *Tubifex*, and *Canthocamptus* (copepod)

Depends on \bar{V}







Types of Palustrine Classes, Subclasses, and Dominance Types


3) Aquatic Bed:

>30% HVC of plants that grow on or below the surface of the water; <30% HVC "taller" plants.

Usually low-energy habitats that are flooded permanently.

- A) Algal: >50% algae *C) Rooted Vascular: >50% RV
 B) Aquatic moss: >50% moss *D) Floating Vascular: >50% FV

Dominance Types: *Chara, Fontinalis, Vallisneria, Ruppia, Nuphar, Lemna, and Eichhornia*



Algal and Aquatic Moss Subclasses

Algae



Spirogyra, Cladophora




Aquatic Moss



Frontinalis spp.
Common Water Moss (streams)

Rooted and Floating Vascular Subclasses



Types of Palustrine Classes, Subclasses, and Dominance Types

4) Unconsolidated Shore:

Shoreline areas with <30% HVC¹ and <75% HC of stone, boulders, or bedrock

Shorelines lacking hydrophytes.

- A) Cobble-gravel: >50% c/g
- B) Sand: >50% sand
- *C) Mud: >50% silt & clay
- D) Organic: >50% dead or live organic matter
- ¹E) Vegetated: >50% non-hydrophytic, pioneer plants that die if flooded.

Dominance Types: Snails, worms, clams, *Xanthium* (cocklebur)



Cobble and Mud Subclasses

Cobble




Mud
"Mudflats"




Organic and Vegetated Subclasses

Organic




Exposed Detritus




Oysters


Vegetated



FACU and UPL Plants



Cocklebur



Types of Palustrine Classes, Subclasses, and Dominance Types


5) Moss-lichen Wetland:


>30% HVC of moss and lichens;
<30% HVC "taller" plants.

Wetlands (bogs and lichen swamps) of northern latitude.

A) Moss: >50% moss B) Lichen: >50% lichen

Dominance Types: Peat moss (*Sphagnum*), liverwort (*Chiloscyphus fragillis*), reindeer moss (*Cladina rangiferina*)





Types of Palustrine Classes, Subclasses, and Dominance Types

6) Emergent Wetland:
 >30% HVC of erect, rooted hydrophytes, excluding mosses and lichens; <30% HVC "taller" plants.

Low energy wetlands (marshes, playas, prairie potholes)


A) **Persistent:** >50% HC of plants that remain standing at least until the beginning of the next growing season.


B) **Nonpersistent:** >50% HC of plants which fall to the surface of the substrate or water at the end of the growing season.

Dominance Types: Cattail (*Typha*), bulrush (*Scirpus*), wild millet (*Echinochloa*), wild rice (*Zizania*), *Panicums*













Types of Palustrine Classes, Subclasses, and Dominance Types

8) Forested Wetland:

Note: Scrub-shrub wetlands can become forested wetlands

>30% HVC of trees (>6m [20 ft.] in height)

Low energy wooded wetlands (swamps, bottoms, hammocks)

- *A) Broad-leaved Deciduous: >50% BLD trees
- *C) Needle-leaved Deciduous: >50% NLD trees
- B) Broad-leaved Evergreen: >50% BLE trees
- D) Needle-leaved Evergreen: >50% NLE trees
- E) Dead: >50% dead trees

Dominance Types: Red maple, green ash, overcup oak, mangroves, baldcypress, tamaracks, white cedar



Forested Wetlands

Broad-leaved Deciduous and Evergreen

BL Deciduous



BL Evergreen



Pneumatophores



Hardwood Bottomlands



Mangrove Forest

Needle-leaved Deciduous and Evergreen

NL Deciduous



NL Evergreen



Cypress Sloughs



Dead Subclass

Scrub-Shrub



Forested



Water Regime Modifiers

(8 Nontidal)

Criteria not as quantitative as USACE zones.

1) Permanently Flooded

Water covers substrate in all years.

2) Intermittently Exposed

Water covers substrate in all years, except during drought.

3) Semi-permanently Flooded

Water covers substrate throughout the growing season in most years.

4) Seasonally Flooded

Water covers substrate for extended periods, especially early in the growing season, but it is absent by the end of the growing season.

Water Regime Modifiers

(8 Nontidal)

Indirect Indicators
 drift & inundation lines, vegetation, etc.

- 5) **Saturated**
 Substrate is saturated, but surface water is seldom present.
- 6) **Temporarily Flooded**
 Surface water is present for brief periods during the growing season, but water table is far below the surface for most of the year.
- 7) **Intermittently Flooded**
 Substrate is usually exposed, but surface water can be present for variable durations but w/o predictable seasonal periodicity.
- 8) **Artificially Flooded**
 Hydroperiod is controlled by pumps, water control structures, and/or levees.

Water Chemistry Modifiers

Salinity:

	PPT
1) Hyperhaline	>40
2) Euhaline	30.0-40
3) Mixohaline	0.5-30
*4) Fresh	<0.5

pH:

	pH
1) Acid	<5.5
2) Circumneutral	5.5-7.4
3) Alkaline	>7.4

Soil Modifiers

Soil Core Depth = 40 cm [16 in] COE

- 1) **Mineral:**

Criteria same as before.

 - A) Saturated Infrequently:
 - <20% dry-weight organic carbon
 - B) Saturated Frequently/Long Durations:
 - <18% dry-weight organic carbon if >60% of mineral portion is clay
 - <12% dry-weight organic carbon if no clay
 - 12-20% dry-weight organic carbon if 0-59% clay
- 2) **Organic:**

<0.002 mm dia.

All substrates that have more organic carbon than above percentages.

Special Modifiers

- 1) **Excavated** Excavated basin or channel.
- 2) **Impounded** Structure (dam) prevents outflow.
- 3) **Diked** Structure (levee) prevents inflow.
- 4) **Partly Drained** Water level has been artificially lowered, but hydrophytes and/or hydric soils present.
- 5) **Farmed** Soil is mechanically disturbed, but hydrophytes will reestablish if farming ceases.
- 6) **Artificial** Non-natural substrate.
e.g., dredge spoil, automobiles, concrete

Alphanumeric Code (Handout)

PSS01-sf04co **PF01-t04cm**
Ω

PEM01-sp04co **PAB03-ie04co**

•1.5 m deep (Sept.)
 •<0.5 ppt; pH = 6.0
 •50% Organic C; 10% Carbon C [trees]

More Practice

PUB02-p04alo **PSS02-ie04alo**
PEM01-sf04alo

PEM02-t04alo

•0.1-1.9 m deep •<0.5 ppt; pH = 8.5
 •Shortly after spring rains •0.1 mm; 30% organic carbon
 •Annuals, perennials, mangrove
